



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## ON A PECULIAR MONSTROSITY IN A FROG.

GEORGE WAGNER,

DEPARTMENT OF ZOÖLOGY, UNIVERSITY OF WISCONSIN.

About three years ago a student, Mr. F. L. Conover, brought me a remarkable specimen of *Rana pipiens*, which he had found among the specimens at the biological laboratory of the Madison High School. The frog was alive. It measured fifty-two millimeters body length and was apparently in full vigor, its death being caused by an overheating of my office at a time when I had to be unexpectedly absent for several days.

The remarkable character of this specimen consists in the presence of three extra limbs extending caudad from the region of the sternum, two on the right and one on the left side. It is very evident that they are attached to an extra basal piece overlying the sternum. All three are more or less imperfect, so that I find it impossible to determine whether they represent arms or legs, or both. Each has four malformed digits, and each shows a thickening indicating the location of the joint between the proximal and distal parts of the limb, the elbow or knee. All are pigmented on the surface away from the frog, with no pigment on the opposite surface. The accompanying photograph (Fig. 2) and radiograph (Fig. 1) of the specimen illustrate better than any verbal description the actual conditions present.

Monstrosities in nature involving the hind legs are not uncommon in the frog, but those involving the fore limbs are apparently rare. The only one known to me is one figured by Sutton ('92, p. 112, Fig. 60), involving the presence of an extra leg on the left side. There is a case in the human being, often quoted, shown in Fig. 3; I have been told about another closely similar case exhibited in this country several years ago, but I have not been able to get exact information concerning it.

Dr. F. R. Lillie kindly called my attention to a very important paper by Tornier (:05) which I would otherwise have overlooked.

Tornier, by an ingeniously chosen cut, separated the upper one quarter or so of the Anlagen of both hind limbs in *Pelobates*. The result was in several cases, at least, the formation of the




FIG. 1.

normal pairs of limbs *plus* two complete extra girdles each with a pair of limbs. Less completely successful experiments resulted in fewer and less fully formed supernumerary parts. But the monstrosities in all cases resembled closely those found in nature, and the close approximation and exposed situation of the Anlagen of the two hind limbs make accidents in nature which may be compared with this artificial "accident" at least not improbable. Tornier's work, it seems to me, gives us a very satisfactory explanation of such forms.

Is it equally applicable to the forelimbs? The location of the Anlagen under the opercle makes experimentation here much more difficult, and no such experiments have been carried out,

so far as I know. One might expect that by operating on one of the Anlagen a supernumerary arm, such as that of Sutton's case, could be produced. It is an experiment that some skilful operator should attempt to perform. But the Anlagen of the two limbs are here not closely approximated; they are far apart. A simultaneous operation on both is not possible, and an accident in nature equally affecting both is in the highest degree improbable.

It appears much more probable to me that in the frog here described we have a case quite parallel to that in Fig. 3, which is



FIG. 2.

properly designated, according to Schwalbe ('07), as a *Thoracopagus parasiticus*. The generally accepted, and to me very probable, explanation of these forms is that they started with a

monstrosity consisting of two complete embryos united at the breast bone. For some reason or other one gains a start in development. As their circulatory systems are united (as is believed, and in many cases demonstrated) this stronger individual (autosite) soon drives its blood into the other (parasite),



FIG. 3. (After Adami, after Wintersohn.)

the heart of which then degenerates through disuse. This means a less adequate blood supply, atrophy, and a lesser or greater absorption by the stronger individual. It is easy to see that the later in the embryonic period the inequality begins the more extensive in development will the parasite be.

I have not attempted to seek all the cases of this sort in man. Approximate parallels of that shown in Fig. 3 must be, to judge from the literature, quite rare. As to the lower vertebrates I have not been able to find a record of any case closely similar to this. It is an interesting fact to note that a frog could grow

to maturity with so very material an impediment. It makes one ask just how much of a variation from the normal is needed to enable natural selection to perform its cruel work?

ZOOLOGICAL LABORATORY,

UNIVERSITY OF WISCONSIN,

July 31, 1913.

#### BIBLIOGRAPHY.

**Adami**

- '10 Principles of Pathology. 2d edition. Vol. 1. Philadelphia, Lea and Febiger.

**Schwalbe**

- '07 Die Morphologie der Missbildungen des Menschen und der Tiere, u.s.w. II. Teil. Jena, Fischer.

**Sutton, J. B.**

- '92 Evolution and Disease. London, Walter Scott.

**Tornier, G.**

- '05 An Knoblauchkröten experimentell entstandene überzählige Hintergliedmassen. Archiv f. Entwickl. Mechanik, 20, pp. 76-124.